10

15

20

WHAT IS CLAIMED IS:

 A method of forming an artificial Shiro of Matsutake comprising:

culturing Matsutake hyphae in a culture substrate containing a substance capable of controlling the cell membrane permeability of the hyphae as an active principle.

A method of forming an artificial Shiro of Matsutake comprising:

culturing Matsutake hyphae in a culture substrate containing a substance capable of enhancing the hydrophilic property of the hyphae as an active principle.

A method of forming an artificial Shiro of Matsutake comprising:

culturing Matsutake hyphae in a culture substrate containing a surfactant and/or a natural vegetable oil as an active principle.

 A method of forming an artificial Shiro of Matsutake comprising:

culturing Matsutake hyphae in a culture substrate containing a fatty acid ester as an active principle.

5. A method of forming an artificial Shiro of Matsutake, comprising:

inducing growth of Matsutake hyphae by aseptically homogenizing a colony of Matsutake hyphae and aseptically culturing the obtained hyphae in a liquid

nutrient medium;

preparing an inoculum of Matsutake hyphae by aseptically replacing the liquid nutrient medium containing the growth-induced Matsutake hyphae with a liquid nutrient medium containing no carbon source; and

culturing aseptically the inoculum of the Matsutake hyphae in a culture substrate containing a substance capable of controlling the cell membrane permeability of the hyphae as an active principle.

A method of forming an artificial Shiro of Matsutake, comprising:

inducing growth of Matsutake hyphae by aseptically homogenizing a colony of Matsutake hyphae and aseptically culturing the obtained hyphae in a liquid nutrient medium;

preparing an inoculum of Matsutake hyphae by aseptically replacing the liquid nutrient medium containing the growth-induced Matsutake hyphae with a liquid nutrient medium containing no carbon source; and

culturing aseptically the inoculum of the Matsutake hyphae in a culture substrate containing a substance capable of enhancing the hydrophilic property of the hyphae as an active principle.

7. A method of forming an artificial Shiro of Matsutake, comprising:

inducing growth of Matsutake hyphae by aseptically homogenizing a colony of Matsutake hyphae and

20

25

15

5

10

15

20

25

aseptically culturing the obtained hyphae in a liquid nutrient medium:

preparing an inoculum of Matsutake hyphae by aseptically replacing the liquid nutrient medium containing the growth-induced Matsutake hyphae with a liquid nutrient medium containing no carbon source; and

culturing aseptically the inoculum of the Matsutake hyphae in a culture substrate containing a surfactant and/or a natural vegetable oil as an active principle.

 A method of forming an artificial Shiro of Matsutake, comprising:

inducing growth of Matsutake hyphae by aseptically homogenizing a colony of Matsutake hyphae and aseptically culturing the obtained hyphae in a liquid nutrient medium;

preparing an inoculum of Matsutake hyphae by aseptically replacing the liquid nutrient medium containing the growth-induced Matsutake hyphae with a liquid nutrient medium containing no carbon source; and

culturing aseptically the inoculum of the Matsutake hyphae in a culture substrate containing a fatty acid ester as an active principle.

9. The method of forming an artificial Shiro of Matsutake according to claim 1, wherein a solution containing the active principle at the concentration of 0.2 to 10 wt% is used as the active principle.

10

15

20

- 10. The method of forming an artificial Shiro of Matsutake according to claim 2, wherein a solution containing the active principle at the concentration of 0.2 to 10 wt% is used as the active principle.
- 11. The method of forming an artificial Shiro of Matsutake according to claim 3, wherein a solution containing the active principle at the concentration of 0.2 to 10 wt% is used as the active principle.
- 12. The method of forming an artificial Shiro of Matsutake according to claim 4, wherein a solution containing the active principle at the concentration of 0.2 to 10 wt% is used as the active principle.
- 13. The method of forming an artificial Shiro of Matsutake according to claim 5, wherein a solution containing the active principle at the concentration of 0.2 to 10 wt% is used as the active principle.
- 14. The method of forming an artificial Shiro of Matsutake according to claim 6, wherein a solution containing the active principle at the concentration of 0.2 to 10 wt% is used as the active principle.
- 15. The method of forming an artificial Shiro of Matsutake according to claim 7, wherein a solution containing the active principle at the concentration of 0.2 to 10 wt% is used as the active principle.
- 16. The method of forming an artificial Shiro of Matsutake according to claim 8, wherein a solution containing the active principle at the concentration of

1.0

15

20

- 0.2 to 10 wt% is used as the active principle.
- 17. The method of forming an artificial Shiro of Matsutake according to claim 1, wherein a solution containing the active principle which is prepared using an organic solvent and distilled water is used as the active principle.
- 18. The method of forming an artificial Shiro of Matsutake according to claim 2, wherein a solution containing the active principle which is prepared using an organic solvent and distilled water is used as the active principle.
- 19. The method of forming an artificial Shiro of Matsutake according to claim 3, wherein a solution containing the active principle which is prepared using an organic solvent and distilled water is used as the active principle.
- 20. The method of forming an artificial Shiro of Matsutake according to claim 4, wherein a solution containing the active principle which is prepared using an organic solvent and distilled water is used as the active principle.
- 21. The method of forming an artificial Shiro of Matsutake according to claim 5, wherein a solution containing the active principle which is prepared using an organic solvent and distilled water is used as the active principle.
 - 22. The method of forming an artificial Shiro of

10

15

20

25

Matsutake according to claim 6, wherein a solution containing the active principle which is prepared using an organic solvent and distilled water is used as the active principle.

- 23. The method of forming an artificial Shiro of Matsutake according to claim 7, wherein a solution containing the active principle which is prepared using an organic solvent and distilled water is used as the active principle.
- 24. The method of forming an artificial Shiro of Matsutake according to claim 8, wherein a solution containing the active principle which is prepared using an organic solvent and distilled water is used as the active principle.
- 25. The method of forming an artificial Shiro of Matsutake according to claim 1, wherein either one of soil having a grain size of 3 mm or less and an artificial substrate having a grain size of 2 mm or less is used as the culture substrate.
- 26. The method of forming an artificial Shiro of Matsutake according to claim 2, wherein either one of soil having a grain size of 3 mm or less and an artificial substrate having a grain size of 2 mm or less is used as the culture substrate.
- 27. The method of forming an artificial Shiro of Matsutake according to claim 3, wherein either one of soil having a grain size of 3 mm or less and

10

15

20

25

an artificial substrate having a grain size of 2 mm or less is used as the culture substrate.

- 28. The method of forming an artificial Shiro of Matsutake according to claim 4, wherein either one of soil having a grain size of 3 mm or less and an artificial substrate having a grain size of 2 mm or less is used as the culture substrate.
- 29. The method of forming an artificial Shiro of Matsutake according to claim 5, wherein either one of soil having a grain size of 3 mm or less and an artificial substrate having a grain size of 2 mm or less is used as the culture substrate.
- 30. The method of forming an artificial Shiro of Matsutake according to claim 6, wherein either one of soil having a grain size of 3 mm or less and an artificial substrate having a grain size of 2 mm or less is used as the culture substrate.
- 31. The method of forming an artificial Shiro of Matsutake according to claim 7, wherein either one of soil having a grain size of 3 mm or less and an artificial substrate having a grain size of 2 mm or less is used as the culture substrate.
- 32. The method of forming an artificial Shiro of Matsutake according to claim 8, wherein either one of soil having a grain size of 3 mm or less and an artificial substrate having a grain size of 2 mm or less is used as the culture substrate.

10

15

20

- 33. The method of forming an artificial Shiro of Matsutake according to claim 1, wherein the active principle is added to the culture substrate in a state of a solution containing the active principle, and weight ratio of the solution containing the active principle to the total weight is 15 to 30 wt%.
- 34. The method of forming an artificial Shiro of Matsutake according to claim 2, wherein the active principle is added to the culture substrate in a state of a solution containing the active principle, and weight ratio of the solution containing the active principle to the total weight is 15 to 30 wt%.
- 35. The method of forming an artificial Shiro of Matsutake according to claim 3, wherein the active principle is added to the culture substrate in a state of a solution containing the active principle, and weight ratio of the solution containing the active principle to the total weight is 15 to 30 wt%.
- 36. The method of forming an artificial Shiro of Matsutake according to claim 4, wherein the active principle is added to the culture substrate in a state of a solution containing the active principle, and weight ratio of the solution containing the active principle to the total weight is 15 to 30 wt%.
- 37. The method of forming an artificial Shiro of Matsutake according to claim 5, wherein the active principle is added to the culture substrate in a state

10

15

20

of a solution containing the active principle, and weight ratio of the solution containing the active principle to the total weight is 15 to 30 wt%.

- 38. The method of forming an artificial Shiro of Matsutake according to claim 6, wherein the active principle is added to the culture substrate in a state of a solution containing the active principle, and weight ratio of the solution containing the active principle to the total weight is 15 to 30 wt%.
- 39. The method of forming an artificial Shiro of Matsutake according to claim 7, wherein the active principle is added to the culture substrate in a state of a solution containing the active principle, and weight ratio of the solution containing the active principle to the total weight is 15 to 30 wt%.
- 40. The method of forming an artificial Shiro of Matsutake according to claim 8, wherein the active principle is added to the culture substrate in a state of a solution containing the active principle, and weight ratio of the solution containing the active principle to the total weight is 15 to 30 wt%.